arXiv:1005.2464v2 [math.FA] 31 Mar 2011

ON HADAMARD TYPE INEQUALITIES INVOLVING SEVERAL KIND OF CONVEXITY

ERHAN SET$^{\star\clubsuit}$, M. EMIN ÖZDEMIR$^{\blacksquare}$, AND SEVER S. DRAGOMIR$^{\blacktriangle}$

Abstract. In this paper, we not only give the extensions of the results given in [7] by Gill et al. for log-convex functions, but also obtain some new Hadamard type inequalities for log-convex, $m-$convex and (α, m)-convex functions.

1. Introduction

The following inequality is well known in the literature as Hadamard's inequality:

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a}\int_a^b f(x)\,dx \leq \frac{f(a)+f(b)}{2}, \tag{1.1}$$

where $f: I \to R$ is a convex function on the interval I of real numbers and $a, b \in I$ with $a < b$. This inequality is one of the most useful inequalities in mathematical analysis. For new proofs, note worthy extension, generalizations and numerous applications on this inequality, see ([1], [4], [5], [6], [9], [12]) where further references are given.

Let I be on interval in R. Then $f: I \to R$ is said to be convex if for all $x, y \in I$ and $\lambda \in [0,1]$,

$$f(\lambda x + (1-\lambda) y) \leq \lambda f(x) + (1-\lambda) f(y)$$

(see [9, P.1]). Geometrically, this means that if K, L and M are three distinct points on the graph of f with L between K and M, then L is on or below chord KM.

Recall that a function $f: I \to (0,\infty)$ is said to be log-convex function, if for all $x, y \in I$ and $t \in [0,1]$, one has the inequality (see [9, P.3])

$$f(tx + (1-t) y) \leq [f(x)]^t [f(y)]^{(1-t)}. \tag{1.2}$$

It is said to be log-concave if the inequality in (1.2) is reversed.

In [13], G. Toader defined m-convexity as follows:

Definition 1. *The function $f: [0,b] \to R$, $b > 0$ is said to be m-convex, where $m \in [0,1]$, if we have*

$$f(tx + m(1-t) y) \leq t f(x) + m(1-t) f(y)$$

for all $x, y \in [0,b]$ and $t \in [0,1]$. We say that f is $m-$concave if $-f$ is $m-$convex.

Denote by $K_m(b)$ the class of all $m-$convex functions on $[0,b]$ for which $f(0) \leq 0$. Obviously, if we choose $m = 1$, definition (1) recaptures the concept of standard convex functions on $[0,b]$.

In [8] , V. G. Miheşan defined $(\alpha, m)-$ convexity as in the following:

2000 *Mathematics Subject Classification.* 26A51, 26D07, 26D15.

Key words and phrases. log-convex functions, $m-$convex functions, $(\alpha, m)-$convex functions, Hadamard's inequality.

*corresponding author.

Definition 2. *The function* $f : [0, b] \to \mathbb{R}$, $b > 0$, *is said to be* $(\alpha, m)-$ *convex, where* $(\alpha, m) \in [0, 1]^2$, *if we have*

$$f(tx + m(1-t)y) \le t^{\alpha} f(x) + m(1-t^{\alpha}) f(y)$$

for all $x, y \in [0, b]$ *and* $t \in [0, 1]$.

Denote by $K_m^{\alpha}(b)$ the class of all $(\alpha, m)-$convex functions on $[0, b]$ for which $f(0) \le 0$. It can be easily seen that for $(\alpha, m) = (1, m)$, $(\alpha, m)-$ convexity reduces to $m-$ convexity and for $(\alpha, m) = (1, 1)$, $(\alpha, m)-$ convexity reduces to the concept of usual convexity defined on $[0, b]$, $b > 0$.

For recent results and generalizations concerning $m-$convex and $(\alpha, m)-$ convex functions, see ([2], [3], [10]).

In [7], P.M. Gill et al. established the following results:

Theorem 1. *Let* f *be a positive,* log-*convex function on* $[a, b]$. *Then*

$$\frac{1}{b-a}\int_a^b f(t)\,dt \le L(f(a), f(b)) \tag{1.3}$$

where

$$L(p, q) = \frac{p-q}{\ln p - \ln q} \quad (p \neq q)$$

is the Logarithmic mean of the positive real numbers p, q *(for* $p = q$, *we put* $L(p, p) = p$*).*

For f *a positive* log-*concave function, the inequality is reversed.*

Corollary 1. *Let* f *be positive* log-*convex functions on* $[a, b]$. *Then*

$$\frac{1}{b-a}\int_a^b f(t)\,dt \le \min_{x\in[a,b]} \frac{(x-a)L(f(a), f(x)) + (b-x)L(f(x), f(b))}{b-a}.$$

If f *is a positive* log-*concave function, then*

$$\frac{1}{b-a}\int_a^b f(x)\,dx \ge \max_{x\in[a,b]} \frac{(x-a)L(f(a), f(x)) + (b-x)L(f(x), f(b))}{b-a}.$$

For some recent results related to the Hadamard's inequalities involving two log-convex functions, see [11] and the references cited therein. The main purpose of this paper is to establish the general version of the inequalities (1.3) and new Hadamard type inequalities involving two log-convex functions or two m-convex functions or two (α, m)-convex functions using elementary analysis.

2. Main Results

We start with the following Theorem.

Theorem 2. *Let* $f_i : I \subset R \to (0, \infty)$ $(i = 1, 2, ..., n)$ *be* log-*convex functions on* I *and* $a, b \in I$ *with* $a < b$. *Then the following inequality holds:*

$$\frac{1}{b-a}\int_a^b \prod_{i=1}^n f_i(x)\,dx \le L\left(\prod_{i=1}^n f_i(a), \prod_{i=1}^n f_i(b)\right) \tag{2.1}$$

where L is a logarithmic mean of positive real numbers.

For f a positive log-*concave function, the inequality is reversed.*

Proof. Since f_i $(i=1,2,...,n)$ are log-convex functions on I, we have

$$f_i\left(ta+(1-t)b\right)\leq\left[f_i\left(a\right)\right]^t\left[f_i\left(b\right)\right]^{(1-t)} \tag{2.2}$$

for all $a,b\in I$ and $t\in[0,1]$. Writing (2.2) for $i=1,2,...,n$, multiplying the resulting inequalities it is easy to observe that

$$\begin{aligned}\prod_{i=1}^{n}f_i\left(ta+(1-t)b\right) &\leq \left[\prod_{i=1}^{n}f_i\left(a\right)\right]^t\left[\prod_{i=1}^{n}f_i\left(b\right)\right]^{(1-t)} \\ &= \prod_{i=1}^{n}f_i\left(b\right)\left[\prod_{i=1}^{n}\frac{f_i\left(a\right)}{f_i\left(b\right)}\right]^t\end{aligned} \tag{2.3}$$

for all $a,b\in I$ and $t\in[0,1]$.

Integrating inequality (2.3) on $[0,1]$ over t, we get

$$\int_0^1\prod_{i=1}^{n}f_i\left(ta+(1-t)b\right)dt\leq\prod_{i=1}^{n}f_i\left(b\right)\int_0^1\left[\prod_{i=1}^{n}\frac{f_i\left(a\right)}{f_i\left(b\right)}\right]^t dt.$$

As

$$\int_0^1\prod_{i=1}^{n}f_i\left(ta+(1-t)b\right)dt=\frac{1}{b-a}\int_a^b\prod_{i=1}^{n}f_i\left(x\right)dx$$

and

$$\int_0^1\left[\prod_{i=1}^{n}\frac{f_i\left(a\right)}{f_i\left(b\right)}\right]^t dt=\frac{1}{\prod\limits_{i=1}^{n}f_i\left(b\right)}L\left(\prod_{i=1}^{n}f_i\left(a\right),\prod_{i=1}^{n}f_i\left(b\right)\right),$$

the theorem is proved. □

Remark 1. *By taking $i=1$ and $f_1=f$ in Theorem 2, we obtain (1.3).*

Corollary 2. *Let $f_i:I\subset R\to(0,\infty)$ $(i=1,2,...,n)$ be* log-*convex functions on I and $a,b\in I$ with $a<b$. Then*

$$\begin{aligned}&\frac{1}{b-a}\int_a^b\prod_{i=1}^{n}f_i\left(x\right)dx \\ &\leq \min_{x\in[a,b]}\frac{(x-a)L\left(\prod\limits_{i=1}^{n}f_i\left(a\right),\prod\limits_{i=1}^{n}f_i\left(x\right)\right)+(b-x)L\left(\prod\limits_{i=1}^{n}f_i\left(x\right),\prod\limits_{i=1}^{n}f_i\left(b\right)\right)}{b-a}.\end{aligned} \tag{2.4}$$

If f_i $(i=1,2,...,n)$ are a positive log-*concave functions, then*

$$\begin{aligned}&\frac{1}{b-a}\int_a^b\prod_{i=1}^{n}f_i\left(x\right)dx \\ &\geq \max_{x\in[a,b]}\frac{(x-a)L\left(\prod\limits_{i=1}^{n}f_i\left(a\right),\prod\limits_{i=1}^{n}f_i\left(x\right)\right)+(b-x)L\left(\prod\limits_{i=1}^{n}f_i\left(x\right),\prod\limits_{i=1}^{n}f_i\left(b\right)\right)}{b-a}.\end{aligned} \tag{2.5}$$

Proof. Let f_i $(i = 1, 2, ..., n)$ be a positive log-convex functions. Then by Theorem 2 we have that

$$\begin{aligned}
&\int_a^b \prod_{i=1}^n f_i(t)\,dt \\
&= \int_a^x \prod_{i=1}^n f_i(t)\,dt + \int_x^b \prod_{i=1}^n f_i(t)\,dt \\
&\leq (x-a)\,L\left(\prod_{i=1}^n f_i(a), \prod_{i=1}^n f_i(x)\right) + (b-x)\,L\left(\prod_{i=1}^n f_i(x), \prod_{i=1}^n f_i(b)\right)
\end{aligned}$$

for all $x \in [a, b]$, whence (2.4). Similarly we can prove (2.5). □

Remark 2. *By taking* $i = 1$ *and* $f_1 = f$ *in (2.4)and (2.5), we obtain the inequalities of Corollary 1.*

We will now point out some new results of the Hadamard type for log-convex, $m-$convex and (α, m)-convex functions, respectively.

Theorem 3. *Let* $f, g : I \to (0, \infty)$ *be* log-*convex functions on* I *and* $a, b \in I$ *with* $a < b$. *Then the following inequalities hold:*

$$\begin{aligned}
&f\left(\frac{a+b}{2}\right) g\left(\frac{a+b}{2}\right) \\
&\leq \frac{1}{2}\left\{\frac{1}{b-a}\int_a^b \left[f(x)\,f(a+b-x) + g(x)\,g(a+b-x)\right]dx\right\} \\
&\leq \frac{f(a)\,f(b) + g(a)\,g(b)}{2}.
\end{aligned} \tag{2.6}$$

Proof. We can write

$$\frac{a+b}{2} = \frac{ta + (1-t)\,b}{2} + \frac{(1-t)\,a + tb}{2}. \tag{2.7}$$

Using the elementary inequality $cd \leq \frac{1}{2}\left[c^2 + d^2\right]$ $(c, d \geq 0$ reals) and equality (2.7), we have

$$\begin{aligned}
&f\left(\frac{a+b}{2}\right) g\left(\frac{a+b}{2}\right) \\
&\leq \frac{1}{2}\left[f^2\left(\frac{a+b}{2}\right) + g^2\left(\frac{a+b}{2}\right)\right] \\
&= \frac{1}{2}\left[f^2\left(\frac{ta+(1-t)\,b}{2} + \frac{(1-t)\,a+tb}{2}\right)\right. \\
&\quad \left. + g^2\left(\frac{ta+(1-t)\,b}{2} + \frac{(1-t)\,a+tb}{2}\right)\right] \\
&\leq \frac{1}{2}\left\{\left[(f(ta+(1-t)\,b))^{\frac{1}{2}}\right]^2 \left[(f((1-t)\,a+tb))^{\frac{1}{2}}\right]^2\right. \\
&\quad \left. + \left[(g(ta+(1-t)\,b))^{\frac{1}{2}}\right]^2 \left[(g((1-t)\,a+tb))^{\frac{1}{2}}\right]^2\right\} \\
&= \frac{1}{2}\left[f(ta+(1-t)\,b)\,f((1-t)\,a+tb)\right. \\
&\quad \left. + g(ta+(1-t)\,b)\,g((1-t)\,a+tb)\right].
\end{aligned} \tag{2.8}$$

Since f, g are log-convex functions, we obtain

$$\begin{aligned} & \frac{1}{2}\left[f\left(ta+(1-t)b\right)f\left((1-t)a+tb\right)\right. \\ & \left.+g\left(ta+(1-t)b\right)g\left((1-t)a+tb\right)\right] \\ \leq\ & \left\{\frac{1}{2}\left[f(a)\right]^{t}\left[f(b)\right]^{(1-t)}\left[f(a)\right]^{(1-t)}\left[f(b)\right]^{t}\right. \\ & \left.+\left[g(a)\right]^{t}\left[g(b)\right]^{(1-t)}\left[g(a)\right]^{(1-t)}\left[g(b)\right]^{t}\right\} \\ =\ & \frac{f(a)f(b)+g(a)g(b)}{2} \end{aligned} \tag{2.9}$$

for all $a, b \in I$ and $t \in [0,1]$.

Rewriting (2.8) and (2.9), we have

$$f\left(\frac{a+b}{2}\right)g\left(\frac{a+b}{2}\right) \leq \frac{1}{2}\left[f\left(ta+(1-t)b\right)f\left((1-t)a+tb\right)+g\left(ta+(1-t)b\right)g\left((1-t)a+tb\right)\right] \tag{2.10}$$

and

$$\begin{aligned} & \frac{1}{2}\left[f\left(ta+(1-t)b\right)f\left((1-t)a+tb\right)+g\left(ta+(1-t)b\right)g\left((1-t)a+tb\right)\right] \\ \leq\ & \frac{f(a)f(b)+g(a)g(b)}{2}. \end{aligned} \tag{2.11}$$

Integrating both sides of (2.10) and (2.11) on $[0,1]$ over t, respectively, we obtain

$$\begin{aligned} & f\left(\frac{a+b}{2}\right)g\left(\frac{a+b}{2}\right) \\ \leq\ & \frac{1}{2}\left[\frac{1}{b-a}\int_a^b\left[f(x)f(a+b-x)+g(x)g(a+b-x)\right]dx\right] \end{aligned} \tag{2.12}$$

and

$$\begin{aligned} & \frac{1}{2}\left[\frac{1}{b-a}\int_a^b\left[f(x)f(a+b-x)+g(x)g(a+b-x)\right]dx\right] \\ \leq\ & \frac{f(a)f(b)+g(a)g(b)}{2}. \end{aligned} \tag{2.13}$$

Combining (2.12) and (2.13), we get the desired inequalities (2.6). The proof is complete. □

Theorem 4. *Let $f, g : I \to (0,\infty)$ be* log-*convex functions on I and $a, b \in I$ with $a < b$. Then the following inequalities hold:*

$$\begin{aligned} & 2f\left(\frac{a+b}{2}\right)g\left(\frac{a+b}{2}\right) \\ \leq\ & \frac{1}{b-a}\int_a^b\left[f^2(x)+g^2(x)\right]dx \\ \leq\ & \frac{f(a)+f(b)}{2}L\left(f(a),f(b)\right)+\frac{g(a)+g(b)}{2}L\left(g(a),g(b)\right) \end{aligned} \tag{2.14}$$

where $L(.\,,\,.)$ is a logarithmic mean of positive real numbers.

Proof. From the inequality (2.10), we have

$$f\left(\frac{a+b}{2}\right)g\left(\frac{a+b}{2}\right) \le \frac{1}{2}\left[f\left(ta+(1-t)b\right)f\left((1-t)a+tb\right) + g\left(ta+(1-t)b\right)g\left((1-t)a+tb\right)\right]$$

for all $a,b\in I$ and $t\in[0,1]$.

Using the elementary inequality $cd\le\frac{1}{2}\left[c^2+d^2\right]$ $(c,d\ge 0$ reals) on the right side of the above inequality, we have

$$f\left(\frac{a+b}{2}\right)g\left(\frac{a+b}{2}\right) \le \frac{1}{4}\left[f^2\left(ta+(1-t)b\right)+f^2\left((1-t)a+tb\right) + g^2\left(ta+(1-t)b\right)+g^2\left((1-t)a+tb\right)\right]. \tag{2.15}$$

Since f,g are log-convex functions, then we get

$$\begin{aligned}
&\left[f^2\left(ta+(1-t)b\right)+f^2\left((1-t)a+tb\right) + g^2\left(ta+(1-t)b\right)+g^2\left((1-t)a+tb\right)\right]\\
\le&\left\{[f(a)]^{2t}[f(b)]^{(2-2t)}+[f(a)]^{(2-2t)}[f(b)]^{2t} + [g(a)]^{2t}[g(b)]^{(2-2t)}+[g(a)]^{(2-2t)}[g(b)]^{2t}\right\}\\
=&\left[f^2(b)\left[\frac{f(a)}{f(b)}\right]^{2t}+f^2(a)\left[\frac{f(b)}{f(a)}\right]^{2t} + g^2(b)\left[\frac{g(a)}{g(b)}\right]^{2t}+g^2(a)\left[\frac{g(b)}{g(a)}\right]^{2t}\right].
\end{aligned}\tag{2.16}$$

Integrating both sides of (2.15) and (2.16) on $[0,1]$ over t, respectively, we obtain

$$2f\left(\frac{a+b}{2}\right)g\left(\frac{a+b}{2}\right)\le\frac{1}{b-a}\int_a^b\left[f^2(x)+g^2(x)\right]dx \tag{2.17}$$

and

$$\begin{aligned}
&\frac{1}{b-a}\int_a^b\left[f^2(x)+g^2(x)\right]dx\\
\le&\frac{1}{2}\left(f^2(b)\int_0^1\left[\frac{f(a)}{f(b)}\right]^{2t}dt+f^2(a)\int_0^1\left[\frac{f(b)}{f(a)}\right]^{2t}dt + g^2(b)\int_0^1\left[\frac{g(a)}{g(b)}\right]^{2t}dt+g^2(a)\int_0^1\left[\frac{g(b)}{g(a)}\right]^{2t}dt\right)\\
=&\frac{1}{2}\left(f^2(b)\left[\frac{\left[\frac{f(a)}{f(b)}\right]^{2t}}{2\log\frac{f(a)}{f(b)}}\right]_0^1+f^2(a)\left[\frac{\left[\frac{f(b)}{f(a)}\right]^{2t}}{2\log\frac{f(b)}{f(a)}}\right]_0^1 + g^2(b)\left[\frac{\left[\frac{g(a)}{g(b)}\right]^{2t}}{2\log\frac{g(a)}{g(b)}}\right]_0^1+g^2(a)\left[\frac{\left[\frac{g(b)}{g(a)}\right]^{2t}}{2\log\frac{g(b)}{g(a)}}\right]_0^1\right)
\end{aligned}\tag{2.18}$$

$$\begin{aligned}
&= \frac{1}{2}\left(\frac{f^2(a)-f^2(b)}{2(\log f(a)-\log f(b))}+\frac{f^2(b)-f^2(a)}{2(\log f(b)-\log f(a))}\right.\\
&\quad\left.+\frac{g^2(a)-g^2(b)}{2(\log g(a)-\log g(b))}+\frac{g^2(b)-g^2(a)}{2(\log g(b)-\log g(a))}\right)\\
&= \frac{1}{2}\left(\frac{f(a)+f(b)}{2}L(f(a),f(b))+\frac{f(a)+f(b)}{2}L(f(b),f(a))\right.\\
&\quad\left.+\frac{g(a)+g(b)}{2}L(g(a),g(b))+\frac{g(a)+g(b)}{2}L(g(b),g(a))\right)\\
&= \left\{\frac{f(a)+f(b)}{2}L(f(a),f(b))+\frac{g(a)+g(b)}{2}L(g(a),g(b))\right\}.
\end{aligned}$$

Combining (2.17) and (2.18), we get the required inequalities (2.14). The proof is complete. □

Theorem 5. *Let $f,g:[0,\infty)\to[0,\infty)$ be such that fg is in $L^1([a,b])$, where $0\le a<b<\infty$. If f is non-increasing m_1-convex function and g is non-increasing m_2-convex function on $[a,b]$ for some fixed $m_1,m_2\in(0,1]$, then the following inequality holds:*

$$\frac{1}{b-a}\int_a^b f(x)g(x)\,dx\le\min\{S_1,S_2\} \tag{2.19}$$

where

$$\begin{aligned}
S_1 &= \frac{1}{6}\left[\left(f^2(a)+g^2(a)\right)+m_1f(a)f\left(\frac{b}{m_1}\right)\right.\\
&\quad\left.+m_2g(a)g\left(\frac{b}{m_2}\right)+m_1^2f^2\left(\frac{b}{m_1}\right)+m_2^2g^2\left(\frac{b}{m_2}\right)\right]
\end{aligned}$$

$$\begin{aligned}
S_2 &= \frac{1}{6}\left[\left(f^2(b)+g^2(b)\right)+m_1f(b)f\left(\frac{a}{m_1}\right)\right.\\
&\quad\left.+m_2g(b)g\left(\frac{a}{m_2}\right)+m_1^2f^2\left(\frac{a}{m_1}\right)+m_2^2g^2\left(\frac{a}{m_2}\right)\right].
\end{aligned}$$

Proof. Since f is m_1-convex function and g is m_2-convex function, we have

$$f(ta+(1-t)b)\le tf(a)+m_1(1-t)f\left(\frac{b}{m_1}\right) \tag{2.20}$$

and

$$g(ta+(1-t)b)\le tg(a)+m_2(1-t)g\left(\frac{b}{m_2}\right) \tag{2.21}$$

for all $t\in[0,1]$. It is easy to observe that

$$\int_a^b f(x)g(x)\,dx=(b-a)\int_0^1 f(ta+(1-t)b)\,g(ta+(1-t)b)\,dt. \tag{2.22}$$

Using the elementary inequality $cd \leq \frac{1}{2}\left(c^2+d^2\right)$ $(c,d\geq 0$ reals$)$, (2.20) and (2.21) on the right side of (2.22) and making the charge of variable and since f,g is non-increasing, we have

$$\begin{aligned}
&\int_a^b f(x)\,g(x)\,dx \\
\leq\ & \frac{1}{2}(b-a)\int_0^1 \left[\{f(ta+(1-t)b)\}^2+\{g(ta+(1-t)b)\}^2\right]dt \\
\leq\ & \frac{1}{2}(b-a)\int_0^1 \left[\left(tf(a)+m_1(1-t)f\left(\frac{b}{m_1}\right)\right)^2\right. \\
& \left.+\left(tg(a)+m_2(1-t)g\left(\frac{b}{m_2}\right)\right)^2\right]dt \\
=\ & \frac{1}{2}(b-a)\left[\frac{1}{3}f^2(a)+\frac{1}{3}m_1^2f^2\left(\frac{b}{m_1}\right)+\frac{1}{3}m_1f(a)f\left(\frac{b}{m_1}\right)\right. \\
& \left.+\frac{1}{3}g^2(a)+\frac{1}{3}m_2^2g^2\left(\frac{b}{m_2}\right)+\frac{1}{3}m_2g(a)g\left(\frac{b}{m_2}\right)\right] \\
=\ & \frac{(b-a)}{6}\left[\left(f^2(a)+g^2(a)\right)+m_1f(a)f\left(\frac{b}{m_1}\right)\right. \\
& \left.+m_2g(a)g\left(\frac{b}{m_2}\right)+m_1^2f^2\left(\frac{b}{m_1}\right)+m_2^2g^2\left(\frac{b}{m_2}\right)\right].
\end{aligned} \tag{2.23}$$

Analogously we obtain

$$\begin{aligned}
&\int_a^b f(x)\,g(x)\,dx \\
\leq\ & \frac{(b-a)}{6}\left[\left(f^2(b)+g^2(b)\right)+m_1f(b)f\left(\frac{a}{m_1}\right)\right. \\
& \left.+m_2g(b)g\left(\frac{a}{m_2}\right)+m_1^2f^2\left(\frac{a}{m_1}\right)+m_2^2g^2\left(\frac{a}{m_2}\right)\right].
\end{aligned} \tag{2.24}$$

Rewriting (2.23) and (2.24), we get the required inequality in (2.19). The proof is complete. □

Theorem 6. *Let $f,g:[0,\infty)\to[0,\infty)$ be such that fg is in $L^1([a,b])$, where $0\leq a<b<\infty$. If f is non-increasing $(\alpha_1,m_1)-$convex function and g is non-increasing $(\alpha_2,m_2)-$convex function on $[a,b]$ for some fixed $\alpha_1,m_1,\alpha_2,m_2\in(0,1]$. Then the following inequality holds:*

$$\frac{1}{b-a}\int_a^b f(x)\,g(x)\,dx \leq \min\{E_1,E_2\} \tag{2.25}$$

where

$$\begin{aligned}
E_1 &= \frac{1}{2}\left[\frac{1}{2\alpha_1+1}f^2(a)+\frac{2\alpha_1^2}{(\alpha_1+1)(2\alpha_1+1)}m_1^2f^2\left(\frac{b}{m_1}\right)\right.\\
&\quad+\frac{2\alpha_1}{(\alpha_1+1)(2\alpha_1+1)}m_1f(a)f\left(\frac{b}{m_1}\right)+\frac{1}{2\alpha_2+1}g^2(a)\\
&\quad+\frac{2\alpha_2^2}{(\alpha_2+1)(2\alpha_2+1)}m_2^2g^2\left(\frac{b}{m_2}\right)\\
&\quad\left.+\frac{2\alpha_2}{(\alpha_2+1)(2\alpha_2+1)}m_2g(a)g\left(\frac{b}{m_2}\right)\right],
\end{aligned}$$

$$\begin{aligned}
E_2 &= \frac{1}{2}\left[\frac{1}{2\alpha_1+1}f^2(b)+\frac{2\alpha_1^2}{(\alpha_1+1)(2\alpha_1+1)}m_1^2f^2\left(\frac{a}{m_1}\right)\right.\\
&\quad+\frac{2\alpha_1}{(\alpha_1+1)(2\alpha_1+1)}m_1f(b)f\left(\frac{a}{m_1}\right)+\frac{1}{2\alpha_2+1}g^2(b)\\
&\quad+\frac{2\alpha_2^2}{(\alpha_2+1)(2\alpha_2+1)}m_2^2g^2\left(\frac{a}{m_2}\right)\\
&\quad\left.+\frac{2\alpha_2}{(\alpha_2+1)(2\alpha_2+1)}m_2g(b)g\left(\frac{a}{m_2}\right)\right].
\end{aligned}$$

Proof. Since f is $(\alpha_1, m_1)-$convex function and g is $(\alpha_2, m_2)-$convex function, then we have

$$f(ta+(1-t)b)\le t^{\alpha_1}f(a)+m_1(1-t^{\alpha_1})f\left(\frac{b}{m_1}\right) \tag{2.26}$$

and

$$g(ta+(1-t)b)\le t^{\alpha_2}g(a)+m_2(1-t^{\alpha_2})g\left(\frac{b}{m_2}\right) \tag{2.27}$$

for all $t\in[0,1]$. It is easy to observe that

$$\int_a^b f(x)g(x)\,dx=(b-a)\int_0^1 f(ta+(1-t)b)g(ta+(1-t)b)\,dt. \tag{2.28}$$

Using the elementary inequality $cd\le\frac{1}{2}\left(c^2+d^2\right)$ ($c,d\ge 0$ reals), (2.26) and (2.27) on the right side of (2.28) and making the charge of variable and since f,g is non-increasing, we have

$$\begin{aligned}
&\int_a^b f(x)g(x)\,dx \\
&\le \frac{1}{2}(b-a)\int_0^1\left[\{f(ta+(1-t)b)\}^2+\{g(ta+(1-t)b)\}^2\right]dt\\
&\le \frac{1}{2}(b-a)\int_0^1\left[\left(t^{\alpha_1}f(a)+m_1(1-t^{\alpha_1})f\left(\frac{b}{m_1}\right)\right)^2\right.\\
&\quad\left.+\left(t^{\alpha_2}g(a)+m_2(1-t^{\alpha_2})g\left(\frac{b}{m_2}\right)\right)^2\right]dt
\end{aligned} \tag{2.29}$$

$$
\begin{aligned}
= & \ \frac{1}{2}(b-a)\left[\frac{1}{2\alpha_1+1}f^2(a)\right. \\
& +\frac{2\alpha_1^2}{(\alpha_1+1)(2\alpha_1+1)}m_1^2 f^2\left(\frac{b}{m_1}\right) \\
& +\frac{2\alpha_1}{(\alpha_1+1)(2\alpha_1+1)}m_1 f(a) f\left(\frac{b}{m_1}\right)+\frac{1}{2\alpha_2+1}g^2(a) \\
& +\frac{2\alpha_2^2}{(\alpha_2+1)(2\alpha_2+1)}m_2^2 g^2\left(\frac{b}{m_2}\right) \\
& \left.+\frac{2\alpha_2}{(\alpha_2+1)(2\alpha_2+1)}m_2 g(a) g\left(\frac{b}{m_2}\right)\right].
\end{aligned}
$$

Analogously we obtain

$$
\begin{aligned}
& \int_a^b f(x) g(x) dx \\
\leq & \ \frac{1}{2}(b-a)\left[\frac{1}{2\alpha_1+1}f^2(b)\right. \\
& +\frac{2\alpha_1^2}{(\alpha_1+1)(2\alpha_1+1)}m_1^2 f^2\left(\frac{a}{m_1}\right) \\
& +\frac{2\alpha_1}{(\alpha_1+1)(2\alpha_1+1)}m_1 f(b) f\left(\frac{a}{m_1}\right)+\frac{1}{2\alpha_2+1}g^2(b) \\
& +\frac{2\alpha_2^2}{(\alpha_2+1)(2\alpha_2+1)}m_2^2 g^2\left(\frac{a}{m_2}\right) \\
& \left.+\frac{2\alpha_2}{(\alpha_2+1)(2\alpha_2+1)}m_2 g(b) g\left(\frac{a}{m_2}\right)\right].
\end{aligned}
\tag{2.30}
$$

Rewriting (2.29) and (2.30), we get the required inequality in (2.25). The proof is complete. □

Remark 3. *In Theorem 6, if we choose $\alpha_1 = \alpha_2 = 1$, we obtain the inequality of Theorem 5.*

REFERENCES

[1] M. Alomari and M. Darus, On the Hadamard's inequality for log-convex functions on the coordinates, *Journal of Inequalities and Applications*, vol. 2009, Article ID 283147, 13 pages, 2009.

[2] M. K. Bakula, M. E. Özdemir and J. Pečarić, Hadamard type inequalities for $m-$convex and $(\alpha, m)-$convex functions, *J. Inequal. Pure & Appl. Math., 9(2008), Article 96.*

[3] M. K. Bakula, J. Pečarić, and M. Ribičić, Companion inequalities to Jensen's inequality for $m-$convex and $(\alpha, m)-$convex functions, *J. Inequal. Pure & Appl. Math., 7(2006), Article 194.*

[4] X.-M. Zhang, Y.-M. Chu and X.-H. Zhang, The Hermite-Hadamard type inequality of GA-convex functions and its application, *Journal of Inequalities and Applications*, vol. 2010, Article ID 507560, 11 pages, 2010.

[5] C. Dinu, Hermite-Hadamard inequality on time scales, *Journal of Inequalities and Applications*, vol. 2008, Article ID 287947, 24 pages, 2008.

[6] S.S. Dragomir and C.E.M. Pearce, Selected Topics on Hermite-Hadamard Inequalities and Applications, *RGMIA Monographs, Victoria University,* 2000. [ONLINE: http://www.staff.vu.edu.au/RGMIA/monographs/hermite-hadamard.html].

[7] P.M. Gill, C.E.M Pearce and J. Pečarić, Hadamard's inequality for r-convex functions, *J. Math. Anal. Appl.*, 215(1997), 461-470.

[8] V. G. Miheşan, A generalization of the convexity, *Seminar on Functional Equations, Approx. and Convex.,* Cluj-Napoca (Romania) (1993)

[9] D.S. Mitrinović, J.E. Pečarić, A.M. Fink, Classical and New Inequalities in Analysis, *Kluwer Academic Publishers, Dordrecht,* 1993.

[10] M.E. Özdemir, M. Avcı and E. Set, On some inequalities of Hermite-Hadamard type via $m-$convexity, *Appl. Math. Lett.* (2010), doi: 10.1016/j.aml.2010.04.037 (in press).

[11] B.G. Pachpatte, A note on integral inequalities involving two *log*-convex functions, *Math. Inequal. Appl.,* 7(4) (2004), 511-515.

[12] E. Set, M.E. Özdemir and S.S. Dragomir, On the Hermite-Hadamard inequality and other integral inequalities involving two functions, *Journal of Inequalities and Applications,* Articles in Press *(2010).*

[13] G. Toader, Some generalizations of the convexity, *Proc. Colloq. Approx. Opt. Cluj-Napoca*, (1984), 329-338.

♣ATATÜRK UNIVERSITY, K.K. EDUCATION FACULTY, DEPARTMENT OF MATHEMATICS, 25240, CAMPUS, ERZURUM, TURKEY
E-mail address: `erhanset@yahoo.com`

■GRADUATE SCHOOL OF NATURAL AND APPLIED SCIENCES, AĞRI İBRAHIM ÇEÇEN UNIVERSITY, AĞRI, TURKEY
E-mail address: `emos@atauni.edu.tr`

▲RESEARCH GROUP IN MATHEMATICAL INEQUALITIES & APPLICATIONS, SCHOOL OF ENGINEERING & SCIENCE, VICTORIA UNIVERSITY, PO BOX 14428, MELBOURNE CITY, MC 8001, AUSTRALIA.
E-mail address: `sever.dragomir@vu.edu.au`
URL: `http://www.staff.vu.edu.au/rgmia/dragomir/`